

## University of Dundee

### Erratum to

Fraser, Callum G.

*Published in:*  
BMC Medicine

*DOI:*  
[10.1186/s12916-016-0713-4](https://doi.org/10.1186/s12916-016-0713-4)

*Publication date:*  
2016

*Licence:*  
CC BY

*Document Version*  
Publisher's PDF, also known as Version of record

[Link to publication in Discovery Research Portal](#)

#### *Citation for published version (APA):*

Fraser, C. G. (2016). Erratum to: Diagnostic work-up of patients presenting in primary care with lower abdominal symptoms: which faecal test and triage strategy should be used? *BMC Medicine*, 14, [158].  
<https://doi.org/10.1186/s12916-016-0713-4>

#### **General rights**

Copyright and moral rights for the publications made accessible in Discovery Research Portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from Discovery Research Portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain.
- You may freely distribute the URL identifying the publication in the public portal.

#### **Take down policy**

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

ERRATUM

Open Access



# Erratum to: Diagnostic work-up of patients presenting in primary care with lower abdominal symptoms: which faecal test and triage strategy should be used?

Callum G. Fraser

## Erratum

After publication of the original article [1], it came to the author's attention that there were two errors in the Conclusions section. 'Quantitative' was mistakenly written as 'qualitative' in the following part of the section:

"**Quantitative** estimates of f-Hb also provide more information than qualitative f-Hb POC tests. Simple and rapid diagnostic pathways are also advantageous. Thus, a possible strategy would be to collect a sample for **quantitative** measure of f-Hb from all patients presenting with lower abdominal symptoms in primary care."

Published online: 12 October 2016

## Reference

1. Fraser CG. Diagnostic work-up of patients presenting in primary care with lower abdominal symptoms: which faecal test and triage strategy should be used? *BMC Med.* 2016;14:139. doi:10.1186/s12916-016-0694-3.

---

Correspondence: callum.fraser@nhs.net  
Centre for Research into Cancer Prevention and Screening, University of Dundee, Ninewells Hospital and Medical School, Dundee DD1 9SY, Scotland, UK



© 2016 The Author(s). **Open Access** This article is distributed under the terms of the Creative Commons Attribution 4.0 International License (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made. The Creative Commons Public Domain Dedication waiver (<http://creativecommons.org/publicdomain/zero/1.0/>) applies to the data made available in this article, unless otherwise stated.